

***In the Claims:***

Please amend claims 70, 72, 73, 75, 80, 82, 90, 92, 93, as shown in the following listing of the entire claims in the Application.

1. - 6. (Canceled)

7. (Withdrawn) A method of producing homozygous stem cells comprising:

(a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by: fusing two oocytes or two spermatids, preventing the extrusion of the second polar body during oogenesis, allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions, or transferring two sperm or two haploid egg nuclei into an enucleated oocyte;

(b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass; and,

(c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

8. (Withdrawn) The method of claim 7, further comprising screening stem cells that are homozygous by genotyping when a mitotically activated post-meiosis I diploid germ cell is produced by (a) fusing two oocytes or two spermatids, or (b) transferring two sperm or two haploid egg nuclei into an enucleated oocyte.

9. (Withdrawn) A method of making a desired progenitor cell, differentiated cell, group of differentiated cells, or tissue type comprising inducing isolated homozygous stem cells to differentiate under suitable conditions.

10. (Withdrawn) The method of claim 9, wherein differentiation is accomplished by the inclusion of a cell regulating factor, hormone or cytokine in the culture medium.

11. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is keratinizing epithelial cells.

12. (Withdrawn) The method of claim 9, wherein said keratinizing epithelial cells are selected from the group consisting of keratinocytes of the epidermis, basal cells of the epidermis, keratinocytes of the fingernails and/or toenails, basal cells of the nail bed, hair shaft cells, hair-root sheath cells, and hair matrix cells.
13. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is cells of wet stratified barrier epithelia.
14. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is epithelial cells specialized for exocrine secretion.
15. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is cells specialized for secretion of hormones.
16. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is epithelial absorptive cells of the gut, exocrine glands, or urogenital tract.
17. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is cells specialized for metabolism and storage.
18. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is epithelial cells serving as the lining the lung, gut, exocrine glands, or urogenital tract or as a barrier.
19. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is epithelial cells lining closed internal body cavities.

20. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is ciliated cells with propulsive function.
21. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is cells specialized for secretion of extracellular matrix.
22. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is contractile cells.
23. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is cells of the blood and immune system.
24. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is sensory transducers.
25. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is autonomic neurons.
26. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is supporting cells of sense organs and of peripheral neurons.
27. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is neurons or glial cells of central nervous system.
28. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is lens cells.

29. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is pigment cells.

30. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is germ cells.

31. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is nurse cells.

32. (Withdrawn) The method of claim 9, wherein the desired cell or group of cells is derived from one of the embryonic germ layers comprising the ectoderm, endoderm or mesoderm.

33. (Withdrawn) A method of producing progenitor cells, comprising:

(a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by: fusing two oocytes or two spermatids, preventing the extrusion of the second polar body during oogenesis, allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions, or transferring two sperm or two haploid egg nuclei into an enucleated oocyte;

(b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass;

(c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass; and,

(d) inducing differentiation of said homozygous stem cells to produce progenitor cells.

34. (Withdrawn) The method of claim 33, further comprising screening stem cells that are homozygous by genotyping when a mitotically activated post-meiosis I diploid germ cell is produced by (a) fusing two oocytes or two spermatids, or (b) transferring two sperm or two haploid egg nuclei into an enucleated oocyte.

35. (Withdrawn) The method of claim 33, further comprising isolating said progenitor cells, and maintaining permanent progenitor cell lines.

36. (Withdrawn) The method of claim 33, wherein cells from said blastocyst-like mass are induced to differentiate in the absence of undifferentiated stem cells.

37. (Withdrawn) A method for producing genetically altered progenitor cells, comprising:

(a) inserting, removing or modifying a desired gene in an HS cell to create a genetically altered HS cell; and,

(b) inducing differentiation of said genetically altered HS cell.

38. (Withdrawn) A method for producing genetically altered progenitor cells, comprising:

(a) inserting, removing or modifying a desired gene in a progenitor cell derived from an HS cell to create a genetically altered progenitor cell; and

(b) culturing said genetically altered progenitor cell to grow genetically altered progenitor cells.

39. (Withdrawn) The methods according to claim 9, 33, or 37, wherein said homozygous stem cells are induced to differentiate in a flat adhesive environment.

40. (Withdrawn) The methods according to claim 9, 33, or 37, wherein said homozygous stem cells are induced to differentiate in a 3D adhesive environment.

41. (Withdrawn) The methods according to claim 9, 33, or 37, wherein said homozygous stem cells are induced to differentiate in a microgravity environment.

42. (Withdrawn) The methods according to claim 9, 33, or 37, wherein said homozygous stem cells are induced to differentiate by generating stemplasms in immunodeficient mice.

43. (Withdrawn) The methods according to claim 33, 37, or 39 wherein said progenitor cells are capable of differentiating into various tissues and cells from only one of the three embryonic layers, the ectoderm, mesoderm, and endoderm.

44. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy.

45. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to treat a disease or condition selected from the group consisting of Parkinson's, Huntington's, Alzheimer's, ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns, foot ulcers, vascular disease, urinary tract disease, AIDS and cancer.

46. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy, wherein the patient is human, and the cells are derived from animal species.

47. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy, wherein the patient is an animal species, and the cells are derived from human.

48. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy, wherein the patient is human, and the cells are derived from human.

49. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy, wherein the patient is an animal species, and the cells are derived animal species.

50. (Withdrawn) A method of therapy comprising administering cells obtained using the methods of claim 9, 33, 37, or 38 to a patient in need of such therapy, wherein the patient is human, and the cells are human.

51. (Withdrawn) The method of claim 50, wherein the cells derived from human are allogenic.

52. (Withdrawn) The method of claim 50, wherein the cells derived from human are isogenic.

53. (Withdrawn) A progenitor cell derived from a method comprising:

- (a) producing a mitotically activated homozygous post-meiosis I diploid germ cell;
- (b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass;
- (c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass; and,
- (d) inducing differentiation of said homozygous stem cells to produce progenitor cells.

54. (Withdrawn) A progenitor cell derived from the method of claim 53, further comprising screening stem cells that are homozygous by genotyping when a mitotically activated post-meiosis I diploid germ cell is produced by (a) fusing two oocytes or two spermatids, or (b) transferring two sperm or two haploid egg nuclei into an enucleated oocyte.

55. (Withdrawn) The progenitor cell of claim 53, further comprising isolating and maintaining said progenitor cells as permanent cell lines.



56. (Withdrawn) The progenitor cell of claim 53, wherein cells from said blastocyst-like mass are induced to differentiate in the absence of undifferentiated stem cells.

57. (Withdrawn) The progenitor cell of claim 53, wherein said homozygous stem cells derived from said blastocyst-like mass are induced to differentiate in a flat adhesive environment.

58. (Withdrawn) The progenitor cell of claim 53, wherein said homozygous stem cells derived from said blastocyst-like mass are induced to differentiate in a 3D adhesive environment.

59. (Withdrawn) The progenitor cell of claim 53, wherein said homozygous stem cells derived from said blastocyst-like mass are induced to differentiate in a microgravity environment.

60. (Withdrawn) The progenitor cell of claim 53, wherein said homozygous stem cells derived from said blastocyst-like mass are induced to differentiate by generating stemplasma in immunodeficient mice.

61. (Withdrawn) The progenitor cell of claim 53, wherein said progenitor cells are capable of differentiating into only one of the three embryonic layers, the ectoderm, endoderm or mesoderm.

62. (Withdrawn) A method of therapy comprising administering the progenitor cells of claim 53 to a patient in need of such therapy.

63. (Withdrawn) A method of therapy comprising administering progenitor cells of claim 53, to treat a disease or condition selected from the group consisting of Parkinson's, Huntington's, Alzheimer's, ALS, spinal cord defects or injuries, multiple sclerosis, muscular dystrophy, cystic

fibrosis, liver disease, diabetes, heart disease, cartilage defects or injuries, burns, foot ulcers, vascular disease, urinary tract disease, AIDS and cancer.

64. (Withdrawn) A method of therapy comprising administering progenitor cells of claim 53 to a patient in need of such therapy, wherein the patient is human, and the differentiated cells are derived from animal species.

65. (Withdrawn) A method of therapy comprising administering progenitor cells of claim 53 to a patient in need of such therapy, wherein the patient is an animal species, and the differentiated cells are derived from human.

66. (Withdrawn) A method of therapy comprising administering progenitor cells of claim 53 to a patient in need of such therapy, wherein the patient is an animal species, and the cells are derived animal species.

67. (Withdrawn) A method of therapy comprising administering progenitor cells of claim 53 to a patient in need of such therapy wherein the patient is human, and the cells are human.

68. (Withdrawn) The method of claim 67, wherein the cells derived from human are allogenic.

69. (Withdrawn) The method of claim 67, wherein the cells derived from human are isogenic.

70. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ obtained from a method comprising:

(a) providing a post-meiosis I diploid oocytes characterized for a given haplotype

- (b) producing a mitotically activated post-meiosis I diploid germ cell homozygous on one or more given haplotypes of ~~said haplotype having an identical pair of one or more alleles~~ by preventing the extrusion of the second polar body during oogenesis or allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (c) culturing said activated post-meiosis I diploid germ cell to form a blastocyst-like mass being surrounded by zona pellucida;
- (d) releasing said blastocyst-like mass from said zona pellucida using the method of assisted hatching; and,
- (e) isolating stem cells homozygous on one or more given haplotypes of ~~said haplotype having an identical pair of one or more alleles~~ from the inner cell mass of said blastocyst-like mass.

71. (previously presented) The parthenogenetically created pluripotent stem cell of claim 70, wherein the method of assisted hatching comprises, fixing said blastocyst-like mass in a micromanipulator having two arms such that one arm holds said blastocyst-like mass in a fixed position, and the other arm applies acidified tyrodes solution to the surface of the zona pellucida in an area equal to about one-eighth of the total surface of said zona pellucida so that the zona pellucida becomes weakened and the blastocyst-like mass is released.

72. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~characterized for a given haplotype having an identical pair of one or more alleles~~ obtained by a method comprising:

- (a) producing a mitotically activated post-meiosis I diploid germ cell homozygous on one or more given haplotypes of ~~a given haplotype having an identical pair of one or more alleles~~ by preventing the extrusion of the second polar body during oogenesis or allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (b) culturing said activated post-meiosis I diploid germ cell to form a blastocyst-like mass being surrounded by zona pellucida;

- (c) transferring said activated post-meiosis I diploid germ cell on day 2 post-activation to a mitomycin C treated feeder layer of mouse embryonic fibroblasts until a blastocyst-like mass is formed;
- (d) releasing said blastocyst-like mass from said zona pellucida using the method of assisted hatching; and,
- (e) isolating stem cells homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ from the inner cell mass of said blastocyst-like mass.

73. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~characterized for a given haplotype having an identical pair of one or more alleles~~ obtained from a method comprising:

- (a) producing a mitotically activated post-meiosis I diploid germ cell homozygous on one or more given haplotypes ~~of said haplotype~~ by allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (b) culturing said activated post-meiosis I diploid germ cell to form a blastocyst-like mass; and,
- (c) isolating stem cells homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ from the inner cell mass of said blastocyst-like mass.

74. (Withdrawn) A method of producing homozygous stem cells comprising:

- (a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass; and,
- (c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

75. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~characterized for a given haplotype having an identical pair of one or more alleles~~ obtained from a method comprising:

- (a) isolating a post-meiosis I diploid germ cell homozygous on one or more given haplotypes ~~of said haplotype~~, wherein said germ cell is an oocyte which matured to metaphase II *in vivo*;
- (b) mitotically activating said metaphase II oocyte by allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (c) culturing said activated metaphase II oocyte to form a blastocyst-like mass; and,
- (d) isolating stem cells homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ from the inner cell mass of said blastocyst-like mass.

76. (Withdrawn) A method of producing homozygous stem cells comprising:

- (a) isolating a homozygous post-meiosis I diploid germ cell, wherein said germ cell is an oocyte which matured to metaphase II *in vivo*;
- (b) mitotically activating said metaphase II oocyte by allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (c) culturing said activated metaphase II oocyte to form a blastocyst-like mass; and,
- (d) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

77. (Withdrawn) A method of making a desired progenitor cell, differentiated cell, group of differentiated cells, or tissue type comprising inducing isolated homozygous stem cells created by the method of claim 73 or 75 to differentiate under suitable conditions.

78. (Withdrawn) An isolated homozygous stem cell derived from a method comprising:

- (a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by preventing the extrusion of the second polar body during oogenesis;

(b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass; and,

(c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

79. (Withdrawn) A method of producing homozygous stem cells comprising:

(a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by preventing the extrusion of the second polar body during oogenesis;

(b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass; and,

(c) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

80. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~characterized for a given haplotype having an identical pair of one or more alleles~~ obtained from a method comprising:

(a) isolating a post-meiosis I diploid germ cell homozygous on one or more given haplotypes of said haplotype, wherein said germ cell is an oocyte which matured to metaphase II *in vivo*;

(b) mitotically activating said metaphase II oocyte by preventing the extrusion of the second polar body during oogenesis;

(c) culturing said activated metaphase II oocyte to form a blastocyst-like mass; and,

(d) isolating stem cells homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ from the inner cell mass of said blastocyst-like mass.

81. (Withdrawn) A method of producing homozygous stem cells comprising:

(a) isolating a homozygous post-meiosis I diploid germ cell, wherein said germ cell is an oocyte which matured to metaphase II *in vivo*;

- (b) mitotically activating said metaphase II oocyte by preventing the extrusion of the second polar body during oogenesis;
- (c) culturing said activated metaphase II oocyte to form a blastocyst-like mass; and,
- (d) isolating homozygous stem cells from the inner cell mass of said blastocyst-like mass.

82. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes ~~characterized for a given haplotype having an identical pair of one or more alleles~~ obtained from a method comprising:

- (a) producing a mitotically activated post-meiosis I diploid germ cell homozygous on one or more given haplotypes ~~of a given haplotype~~ by preventing the extrusion of the second polar body during oogenesis or allowing the extrusion of the second polar body and spontaneous self-replication under appropriate conditions;
- (b) culturing said activated post-meiosis I diploid germ cell to form a blastocyst-like mass;
- (c) transplanting said blastocyst-like mass into an animal host to create a stemplasm; and,
- (d) isolating stem cells homozygous on one or more given haplotypes ~~of a given haplotype having an identical pair of one or more alleles~~ from said stemplasm.

83. (Canceled)

84. (Previously presented) The parthenogenetically created stem cell of claim 82, wherein said blastocyst-like mass is transplanted under the kidney capsule.

85. (Withdrawn) A method of producing homozygous stem cells comprising:

- (a) producing a mitotically activated homozygous post-meiosis I diploid germ cell by: fusing two oocytes or two spermatids, preventing the extrusion of the second polar body during oogenesis, allowing the extrusion of the second polar body and spontaneous self-replication

under appropriate conditions, or transferring two sperm or two haploid egg nuclei into an enucleated oocyte;

(b) culturing said activated homozygous post-meiosis I diploid germ cell to form a blastocyst-like mass;

(c) transplanting said blastocyst-like mass into an animal host to create a stemplasm; and,

(d) isolating homozygous stem cells from said stemplasm.

86. (Withdrawn) The method of claim 85, further comprising screening stem cells that are homozygous by genotyping when a mitotically activated post-meiosis I diploid germ cell is produced by (a) fusing two oocytes or two spermatids, or (b) transferring two sperm or two haploid egg nuclei into an enucleated oocyte.

87. (Withdrawn) The method of claim 85, wherein said blastocyst-like mass is transplanted under the kidney capsule.

88. (Withdrawn) A method of making a desired progenitor cell, differentiated cell, group of differentiated cells, or tissue type comprising inducing isolated homozygous stem cells created by the method of claim 78, 80, 82, or 85 to differentiate under suitable conditions.

89. (Canceled)

90. (Currently Amended) A parthenogenetically created pluripotent stem cell homozygous on one or more given haplotypes. ~~characterized for a given haplotype having an identical pair of one or more alleles.~~

91. (Canceled)



92. (Currently amended) The parthenogenetically created stem cell of claim 70, 71, 72, 73, 75, 80, 82, 84, or 90 homozygous on one or more given haplotypes ~~having a given haplotype with an identical pair of one or more alleles~~ wherein said stem cell is obtained from human.

93. (Currently amended) The parthenogenetically created stem cell of claim 70, 71, 72, 73, 75, 80, 82, 84, or 90 homozygous on one or more given haplotypes ~~having a given haplotype with an identical pair of one or more alleles~~ wherein said stem cell is obtained from non-human.

94. (Previously presented) The parthenogenetically created stem cell of claim 93, wherein the non-human species is selected from the group consisting of mouse, hamster, dog, cat, rabbit, ferret, mink, guinea pig, hedgehogs, cattle, sheep, goat, llama, horse, deer, pig, monkey, and ape.

95. (Previously presented) The parthenogenetically created stem cell of claim 92 wherein the stem cell contains an identical set of Major Histocompatibility alleles.